


**SYMPOSIUM ON  
COMBINATORICS AND GRAPH THEORY**

*25 February — 29 February, 1980*

NEW INEQUALITIES FOR THE PARAMETERS OF  
AN ASSOCIATION SCHEME

A. NEUMAIER

Technische Universität Berlin



INDIAN STATISTICAL INSTITUTE  
203 BARRACKPORE TRUNK ROAD  
CALCUTTA 700 035

## New inequalities for the parameters of an association scheme

by A. Neumaier, Fachbereich Mathematik,  
Technische Universität Berlin, D-1000 Berlin 12

Let  $X$  be a finite set with  $v$  objects. An  $s$ -class scheme on  $X$  is a partition of the set of all 2-subsets of  $X$  into  $s \geq 2$  nonempty classes. Two objects  $x$  and  $y$  are  $i$ -th associates if  $x \neq y$  and  $\{x, y\}$  is in the  $i$ -th class of the partition. We define  $k_i(x)$  as the number of  $i$ -th associates of  $x$ , and  $p_{ij}(x, y)$  as the number of  $z \in X$  which are  $i$ -th associates of  $x$ , and  $j$ -th associates of  $y$ . If we write  $D_i = (d_{xy}^i)$  with  $d_{xy}^i = 1$  or  $0$  according as  $x$  and  $y$  are  $i$ -th associates or not then

$$D_i D_j = (p_{ij}(x, y)), \quad (1)$$

$$p_{ij}(x, x) = k_i(x) \delta_{ij}, \quad (1')$$

where  $\delta_{ij}$  ( $= 1$  or  $0$  according as  $i=j$  or not) is the Kronecker symbol. An association scheme is a scheme with  $k_i(x) = k_i$  for all  $x \in X$ , and  $p_{ij}(x, y) = p_{ij}^e$  whenever  $x$  and  $y$  are  $\ell$ -th associates.

For the following well-known results see e.g. Cameron, et al [1].

The Bose-Mesner-algebra of an association scheme is the algebra  $V$  generated by the matrices  $I$  and  $D_1, \dots, D_s$ . There is a basis  $E_0, \dots, E_s$  of  $V$  satisfying for  $i, j = 0, \dots, s$

$$E_i E_j = \delta_{ij} E_j. \quad (2)$$

Also, the Bose-Mesner-algebra is closed under pointwise multiplication  $(a_{xy}) \circ (b_{xy}) = (a_{xy} b_{xy})$ , whence

$$E_i \circ E_j = \sum_{\ell=0}^s q_{ij}^{\ell} E_{\ell} \quad (3)$$

for appropriate numbers  $q_{ij}^{\ell}$ , which can be calculated from the parameters. The Krein condition

$$q_{ij}^{\ell} \geq 0 \text{ for } i, j, \ell = 0, \dots, s \quad (4)$$

gives a restriction on the parameters.

The ranks  $f_i$  of  $E_i$  can also be calculated from the parameters; they satisfy  $f_0 + \dots + f_s = v$ , and the fact that all  $f_i$  must be integers places a severe restriction on the possible parameter sets. We prove here a new inequality for the ranks:

Theorem 1

The following inequalities hold:

$$\sum_{\ell: q_{ij}^{\ell} > 0} f_{\ell} < \begin{cases} f_i f_j & \text{for } i \neq j, \\ \frac{1}{2} f_i (f_i + 1) & \text{for } i = j. \end{cases} \quad (5)$$

Proof. By the following lemma, the rank of  $E_i \circ E_j$  is at most  $f_i f_j$  if  $i \neq j$ , and  $\frac{1}{2} f_i (f_i + 1)$  if  $i = j$ . On the other hand, since the  $E_i$  are mutually orthogonal, (3) implies that the rank of  $E_i \circ E_j$  is given by the left hand side of (5).

Lemma

- (i) Let  $A$  be a matrix of rank  $f$ . Then  $A \circ A$  has rank  $\leq \frac{1}{2} f(f+1)$ .
- (ii) Let  $A$  and  $B$  be matrices of the same size of rank  $f$  resp.  $g$ . Then  $A \circ B$  has rank  $\leq fg$ .

Proof. Write  $A = (a_{xy})$ , and let  $x_1, \dots, x_f$  be the labels of  $f$  independent rows. Then each  $a_{xy}$  is a linear combination of  $a_{x_1y}, \dots, a_{x_fy}$ . Hence each entry  $a_{xy}^2$  of  $A \circ A$  is a linear combination of  $a_{x_1y}^2, \dots, a_{x_fy}^2 + a_{x_1y}a_{x_2y}, \dots, a_{x_{f-1}y}a_{x_fy}$ , of which there are  $f + \binom{f}{2} = \frac{1}{2}f(f+1)$  terms. This proves (i), and the proof of (ii) is completely analogously.

A 2-class association scheme is essentially the same as a strongly regular graph (see e.g. Seidel [2] for a definition). A strongly regular graph with  $q_{22}^2 = 0$  is called a Smith graph. Cameron, et al [1] show that  $q_{22}^0$  and  $q_{22}^1$  are nonzero. Hence theorem 1 gives

Theorem 2

(i) The parameters of a strongly regular graph which is not a Smith graph satisfy

$$v \leq \frac{1}{2}f_2(f_2+1). \tag{6}$$

(ii) The parameters of a Smith graph satisfy

$$v \leq \frac{1}{2}f_2(f_2+3). \tag{7}$$

Proof. Apply theorem 1 with  $i=j=2$ , and observe that  $f_0+f_1+f_2 = v$ .

Example

The following parameter set for a strongly regular graph satisfies all previously known conditions for strongly regular graphs (as stated e.g. in Seidel [2]) but fails (6):

$$v=841, k=200, \lambda=87, \mu=35, f_2=40.$$

Problem. Characterize those graphs for which (6) is satisfied with equality.

Remarks. 1. More inequalities can be obtained similarly by looking at  $E_i \circ E_j \circ E_k$ , etc., but it is not known whether they are really more restrictive than those of theorems 1 and 2.

2. The special case of theorem 2, where the graph has a rank 3 automorphism group, has been proved already in Cameron, et al [1].

3. Theorem 2 improves the absolute bound (see e.g. Seidel [2]) for strongly regular graphs; it is not known how theorem 1 relates to the more general absolute bound mentioned in [1], proposition 6.1.

### References

1. P.J. Cameron, J.M. Goethals, and J.J. Seidel, The Krein condition, spherical designs, Norton algebras and permutation groups, Proc. Kon. Ned. Acad. Wet. A81 (1978), 196-206.
2. J.J. Seidel, Strongly regular graphs, an introduction, Proc. 7<sup>th</sup> Brit. Comb. Conf., Cambridge (LMS Lecture Notes Series 38) 1979, pp. 157-180.