

Some Relations between Roots, Holes, and Pillars

A. NEUMAIER

These are some observations originating in work by Conway and Sloane (*Proc. R. Soc. London* **A381** (1982), 275-283) on holes of the Leech lattice and by Lemmens and Seidel (*J. Algebra* **24** (1973), 494-512) on equiangular lines. We observe close relations which may be relevant for applications to the classification of maximal twographs with smallest eigenvalue -5 , and of integral lattices generated by norm 4 vectors.

1. ROOT CLASSES

Let L be an integral even lattice of minimum norm 4 and dimension n . On the set of all norm 2 vectors of \mathbb{R}^n we define an equivalence relation $r \equiv s$ iff $r - s \in L$. For reasons explained in a moment we call the equivalence classes R of \equiv the *root classes* of L . We are interested in *rich root classes* R , i.e. root classes R such that $|R| \geq n + 1$ and R spans \mathbb{R}^n , in the hope that their structure is sufficient to reconstruct L . Let R be a root class. Clearly

$$r - s \in L \text{ for all } r, s \in R, \tag{1.1}$$

$$r \in R + L, \quad (r, r) = 2 \Rightarrow r \in R. \tag{1.2}$$

For $r, s \in R, r \neq s$ we have $(r, s)^2 \leq (r, r)(s, s) = 4$ and

$$4 \leq (r - s, r - s) = (r, r) - 2(r, s) + (s, s) = 4 - 2(r, s) \tag{1.3}$$

whence

$$(r, s) = 2 - \frac{1}{2}(r - s, r - s) \in \{0, -1, -2\}. \tag{1.4}$$

In particular, $(r, s) \leq 0$ for $r, s \in R, r \neq s$ so that R is a (reduced or extended) fundamental system of roots of equal length 2.

We denote by h_R the greatest common divisor of the set $\{\sum c_r | c_r \in \mathbb{Z}, \sum c_r r = \theta\}$, where R is the index set. We then have

$$\sum c_r r = \theta, c_r \in \mathbb{Z} \Rightarrow \sum c_r \equiv 0 \pmod{h_R}, \tag{1.5}$$

$$h_R R \subseteq L; \tag{1.6}$$

the last relation (1.6) follows since $(\sum c_r)s = \sum c_r(s - r) \in L$ for all $s \in R$. Note that h_R is zero if and only if the roots in R are linearly independent; in particular $h_R \neq 0$ if R is a rich root class.

We remark that in the above setting, if we drop the assumption of minimum norm 4 for a root class R and $r, s \in R$ we only have $(r, s) \in \{0, \pm 1, \pm 2\}$ in place of (1.4); therefore R is still a set of roots (consider the root system defined by the set of norm 2 vectors of the integral lattice generated by R), but R is no longer necessarily a fundamental system.

EXAMPLE. Let $n = 24$, and let $L = A_{24}$ be the Leech lattice. If r is a deepest hole nearest to the origin (Conway and Sloane [3]) then $(r, r) = 2$ and the root class R containing r is a fundamental system of one of the types $\hat{A}_1^{24}, \hat{A}_2^{12}, \hat{A}_3^8, \dots, \hat{D}_{16}\hat{E}_8, \hat{D}_{24}$, depending on the type of the hole. In each case the root class is rich; moreover, L can be reconstructed from R by one of the 23 constructions described in Conway and Sloane [4], cf. also Borchers [1].

2. WEYL VECTORS

From the discussion of the 23 constructions in [4] (where a reference to Bourbaki [2] is given), we take the following properties of fundamental systems.

Let \hat{R} be an irreducible extended fundamental system. The adjacency matrix of the Dynkin diagram of \hat{R} has a positive integral Perron vector c (corresponding to the eigenvalue 2) whose smallest component is 1; clearly

$$\sum c_r r = 0. \quad (2.1)$$

The number $h = \sum c_r$ is called the *Coxeter number* of \hat{R} , and $n = |\hat{R}| - 1$ is the dimension of the lattice $L(\hat{R})$ spanned by \hat{R} . The number of roots $r_x \in \hat{R}$ with $c_r = 1$ (which we call the *extremal roots* of \hat{R}) equals the determinant of $L(\hat{R})$, the order of $L(\hat{R})^*/L(\hat{R})$, and the order of the symmetry group of the Dynkin diagram of \hat{R} . For an extremal root r_x of \hat{R} , the set $\hat{R} \setminus \{r_x\}$ is a reduced fundamental system of roots, and there is a unique vector $w_x \in \langle \hat{R} \rangle$, the *Weyl vector* of $\hat{R} \setminus \{r_x\}$, such that

$$(r, w_x) = \begin{cases} h - 1, & \text{for } r = r_x, \\ -1, & \text{for } r \in \hat{R} \setminus \{r_x\}; \end{cases} \quad (2.2)$$

the Weyl vector can alternatively be described as half the sum of the positive roots of the root system defined by $\hat{R} \setminus \{r_x\}$. The norm of the Weyl vector is

$$(w_x, w_x) = \frac{n}{12} h(h + 1). \quad (2.3)$$

One easily sees that if r_x and r_y are distinct extremal roots of \hat{R} then

$$\frac{1}{h} (w_x - w_y) \in L(\hat{R})^*. \quad (2.4)$$

For example (with orthonormal vectors e_i), if $\hat{R} = \hat{A}_n = \{e_i - e_{i+1} | i \bmod n + 1\}$ and $r_x = e_{n+1} - e_1$ we have

$$h = n + 1, \quad w_x = \frac{1}{2} \sum_{j=1}^{n+1} (n + 2 - 2j) e_j \quad \text{of norm } \frac{n(n+1)(n+2)}{12}, \quad (2.5)$$

and if $\hat{R} = \hat{D}_n = \{e_i - e_{i+1} | i = 1, \dots, n-1\} \cup \{-e_1 - e_2, e_{n-1} + e_n\}$ and $r_x = -e_1 - e_2$ we have

$$h = 2n - 2, \quad w_x = \sum_{j=1}^n (n - j) e_j \quad \text{of norm } \frac{n(n-1)(2n-1)}{6}. \quad (2.6)$$

Note that Conway and Sloane [4] consider instead the scaled Weyl vectors $g_x = -(1/h)w_x$.

3. GENERALIZING THE 23 CONSTRUCTIONS

We are interested in the question whether L can be reconstructed from the sublattice

$$L_R = \left\{ \sum m_r r \mid m_r \in \mathbb{Z}, \sum m_r = 0 \right\} = \left\{ \sum m_r r \mid \sum m_r \equiv 0 \pmod{h_R} \right\} \quad (3.1)$$

generated by the differences of roots of a rich root class R . Since R is assumed rich, $h_R \neq 0$, and the dimensions of L_R and L agree. To reconstruct L we have to find the remaining vectors $z \in L \setminus L_R$. Since L is integral, $z \in L_R^*$; in particular $(z, r - s) \in \mathbb{Z}$ for $r, s \in R$, and $(z, h_R r) \in \mathbb{Z}$. Hence there is an integer α with

$$(z, r) \equiv -\frac{\alpha}{h_R} \pmod{1}, \quad \text{for all } r \in R. \quad (3.2)$$

Therefore, for the projection z_i of z to the subspace spanned by an irreducible component R_i of R we have

$$\left(z_i - \frac{\alpha w_x}{h_R}, r \right) \in \mathbb{Z}, \quad \text{for } r \in R_i, \tag{3.3}$$

where x is an extremal node of R_i (if R_i is extended) or of \hat{R}_i (if R_i is reduced) and w_x is the corresponding Weyl vector. This implies

$$z_i \in \alpha w_i + L(R_i)^*. \tag{3.4}$$

Note that if R_i is extended then, by construction of R_i , the number h_R divides the Coxeter number h_i of R_i . But since R is rich, at least one R_i is extended, so that there are only finitely many possibilities for the z_i and hence for $z = \sum z_i$. The requirement that z has integral norm further restricts the possibilities for the z_i . To find the (only finitely many) lattices admitting a given fundamental system R as a rich root class one then has to check when two vectors with given components z_i (according to (3.4)) have integral inner product. Knowing this it is easy to construct the glue code $L/L_R \subseteq L_R^*/L_R$. Although this will involve considerable work, it is a finite process for each R . Preliminary calculations of the author suggest the following.

CONJECTURE. (2.3) and (3.2)–(3.4) imply that $n = \sum n_i < 48$.

The truth of the conjecture would make it feasible to find all lattices with a rich root class.

In principle the above analysis can be extended to all root classes R with $h_R \neq 0$; however then only the projection of L to the subspace spanned by L_R can be determined and further analysis is needed to reconstruct L .

4. EQUIANGULAR LINES WITH ANGLE ARCCOS 1/5

Let L be an even lattice, and let $L^{(m)} = \{x \in L \mid (x, x) = m\}$. Fix $e \in L^{(6)}$, and put

$$X := \left\{ x - \frac{e}{2} \mid x \in L^{(4)}, (x, e) = 3 \right\}.$$

Then X is a set of vectors of norm $5/2$. Since for $x, y \in X$, $(x - y, x - y)$ and $(x + y - e, x + y - e)$ are nonnegative even numbers, vectors in X have mutual inner products in $\{\pm 1/2, \pm 3/2, \pm 5/2\}$; the lines along these vectors therefore have mutual angles arccos $1/5$ or arccos $3/5$. Moreover if L contains no norm 2 vectors then the inner products $\pm 3/2$ cannot occur since $(x - e/2, y - e/2) = \pm 3/2$ implies $x - y \in L^{(2)}$ for the upper sign and $x + y - e \in L^{(2)}$ for the lower sign; therefore, in this case X defines a set of equiangular lines with angles arccos $1/5$. Such sets have been studied by Lemmens and Seidel [6] and are in close relation with twographs whose smallest eigenvalue is -5 (Taylor [10]). Conversely, if X is a set of vectors of norm $5/2$ along lines at angles arccos $1/5$ or arccos $3/5$ then, with a vector $e \perp X$ of norm 6, the vectors $e/2 \pm x$ ($x \in X$) have norm 4 and integral inner product and hence span an even lattice L . However, even if X is equiangular with angles arccos $1/5$, it is not clear whether L has no norm 2 vectors.

Now suppose that X contains a 6-clique, i.e. six vectors $x_1, \dots, x_6 \in X$ with mutual inner product $-1/2$. Put $u = (x_1 + x_2 + x_3)/3$ and $v = (x_4 + x_5 + x_6)/3$. Then $(u, u) = -(u, v) = (v, v) = 1/2$; in particular $v = -u$. Using $(x_i, u) = 1/2$ for $i \leq 3$ and $(x_i, u) = -1/2$ for $i \geq 4$, it is easily verified that the vectors $r_i = u - x_i$ ($i = 1, 2, 3$), $s_i = u + x_{i+3}$ ($i = 1, 2, 3$), and $t_1 = e/2 + 4$, $t_2 = u - e/2$, $t_3 = -2u$ form an extended root system of type \tilde{A}_2^3 . Moreover, one easily sees that the 9 roots r_i, s_i, t_i ($i = 1, 2, 3$) belong to the same root class R of L . The remaining members of R are the roots $u \pm x$, where x

belongs to the set

$$P := \{x \in X \mid (x, x_i) = 1 \text{ for } i \leq 3, (x, x_i) = -1 \text{ for } i \geq 4\}.$$

In the terminology of Lemmens and Seidel [6], P is the *pillar* determined by the sign distribution $\varepsilon = (+++---)$ on the 6-clique X . This explains the occurrence of Dynkin diagrams in the pillar analysis of Lemmens and Seidel. As an example, the pillars of the 276 equiangular lines in \mathbb{R}^{23} constructed by Taylor [10] from the Leech lattice consist of 9 triangles, giving the root class $R = \hat{A}_2^{12}$ and the Leech lattice $L = A_{24}$. This fact is also related to the uniqueness proof of the corresponding regular twograph by Goethals and Seidel [5].

5. HOLES

It remains to consider the question how to find rich root classes of a lattice L . In view of the examples from the Leech lattice it is natural to consider holes.

Let L be an even lattice of dimension n and minimal norm 4, and let f be a hole of radius $R = \sqrt{m}$ nearest to the origin. Then $(f, f) = m$. We call the hole *spherical*, *affine*, or *hyperbolic*, depending on whether $m < 2$, $m = 2$, $m > 2$. For example, by Conway and Sloane [3], all holes of the Leech lattice are spherical (small holes) or affine (deepest holes). Since f is a hole, the set Σ of nearest lattice points has $s + 1 \geq n + 1$ points $v_i = f - f_i$ ($i = 0, \dots, s$), say; thus $f = f_0$ and

$$(f_i, f_i) = m. \quad (5.1)$$

Since f is a hole, f is a strictly convex combination of the v_i , $f = \sum c_i(f - f_i)$, $c_i > 0$, $\sum c_i = 1$, so that

$$\sum c_i f_i = 0 \quad \text{with } c_i > 0, \quad \sum c_i = 1; \quad (5.2)$$

moreover the v_i and hence the f_i span \mathbb{R}^n . In particular, if $m = 2$ then $\{f_0, \dots, f_s\}$ is a rich root class.

Now (5.1) implies

$$(f_i, f_j) = m - \frac{1}{2}(f_i - f_j, f_i - f_j); \quad (5.3)$$

therefore, for $i \neq j$, $(f_i, f_j) = m - p$ with an integer $p \geq 2$. This implies that the vectors f_0, \dots, f_s have a Gram matrix of the form

$$G = 2I - A + (m - 2)J, \quad (5.4)$$

where $A = (a_{ij})$ is an integral nonnegative matrix and J is the matrix all of whose entries are 1. This allows us to define a multigraph Γ with vertices f_i , where for $i \neq j$, f_i and f_j are joined by a_{ij} edges, and $(f_i, f_j) = m - 2 - a_{ij}$. Clearly G is positive semidefinite of rank n , and by (5.2), $c = (c_0, \dots, c_s)^T$ is a positive null vector of G ; i.e. with $j = (1, \dots, 1)^T$, we have

$$Gc = 0, \quad c > 0, \quad j^T c = 1. \quad (5.5)$$

We now distinguish several cases.

(a) For a spherical hole we have $m < 2$. Then $2I - A$ is positive definite whence A has largest eigenvalue < 2 . Therefore, Γ is a disjoint union of spherical Dynkin diagrams.

(b) For an affine hole we have $m = 2$. Then $2I - A$ is positive semidefinite, and (5.5) implies $Ac = 2c > 0$. Therefore Γ is a disjoint union of affine Dynkin diagrams.

(c) For a hyperbolic hole we have $m > 2$. Then $Ac > 2c > 0$ whence the largest eigenvalue $\lambda_1(A)$ of A is > 2 . But since G is positive semidefinite and J has rank 1, $2I - A = G - (m - 2)J$ has inertia $(-1)(+1)^{n-1}0^{s+1-n}$ or $(-1)(+1)^n0^{s-n}$. Therefore, the second largest eigenvalue $\lambda_2(A)$ of A is ≤ 2 ,

$$\lambda_1(A) > 2 \geq \lambda_2(A). \quad (5.6)$$

In particular, Γ is connected. Since $2\mathbf{I} - \mathbf{A}$ may be regarded as the Gram matrix of a set of norm 2 vectors in hyperbolic space $\mathbb{R}^{n,1}$ we are justified in calling a multigraph Γ whose adjacency matrix \mathbf{A} satisfies (5.6) a *hyperbolic multigraph*; cf. e.g. Maxwell [7], Neumaier [8]. From (5.5) we find the further relations

$$(\mathbf{A} - 2\mathbf{I})\mathbf{c} = (m - 2)\mathbf{j}, \quad c > 0, \quad (5.7)$$

$$m = 2 + 1/\mathbf{j}^T\mathbf{c}. \quad (5.8)$$

The fact that (5.7) must have a positive solution restricts the possibilities for Γ a little; however, e.g. all regular hyperbolic multigraphs with v points and valency k might still occur with $m = 2 + (k - 2)/v$. In particular, many sporadic strongly regular graphs, like the Hoffman–Singleton graph ($v = 50$, $m = 21/10$), the Higman–Sims graph ($v = 100$, $m = 11/5$) and the McLaughlin graph ($v = 275$, $m = 12/5$) are candidates for Γ ; cf. [5], [9].

REFERENCES

1. R. E. Borcherds, The Leech lattice, *Proc. R. Soc. London* **A398** (1985) 365–376.
2. N. Bourbaki, *Groupes et algèbres de Lie*, Hermann, Paris, 1968, ch. IV, V, VI.
3. J. H. Conway and N. J. A. Sloane, The covering radius of the Leech lattice, *Proc. R. Soc. London* **A380** (1982) 261–290.
4. J. H. Conway and N. J. A. Sloane, Twenty-three constructions for the Leech lattice, *Proc. R. Soc. London* **A381** (1982) 275–283.
5. J. M. Goethals and J. J. Seidel, The regular two-graph on 276 vertices, *Discrete Math.* **12** (1975), 143–158.
6. P. W. H. Lemmens and J. J. Seidel, Equiangular lines, *J. Algebra* **24** (1973), 494–512.
7. G. Maxwell, Hyperbolic trees, *J. Algebra* **54** (1978) 46–49.
8. A. Neumaier, The second largest eigenvalue of a tree, *Linear Algebra Appl.* **46** (1982), 9–25.
9. A. Neumaier and J. J. Seidel, Discrete hyperbolic geometry, *Combinatorica* **3** (1983), 219–237.
10. D. E. Taylor, Regular 2-graphs, *Proc. London Math. Soc.* **35** (1977), 257–274.

Received 13 June 1985 and in revised form 12 November 1985

A. NEUMAIER

Institut für Angewandte Mathematik, Universität Freiburg,
D-7800 Freiburg i. Br., F.R.G.